



Drug (s)	Duzallo (lesinurad/allopurinol Zurampic (lesinurad)
Policy#	14101
Indications	Zurampic is indicated for the adjunct treatment of hyperuricemia associated with gout (in combination with a xanthine oxidase inhibitor, ie, Allopurinol or Uloric) in patients who have not achieved target serum uric acid levels with a xanthine oxidase inhibitor alone.  Duzallo is indicated for the treatment of hyperuricemia associated with gout in patients not achieving target serum uric acid levels with allopurinol alone.
CRITERIA	ConnectiCare considers <b>Zurampic and Duzallo</b> to be medically necessary in adults who meet all of the following criteria:  • Patient has clinically diagnosed hyperuricemia associated with gout • Patient has not achieved a target serum uric acid level using allopurinol (Zyloprim) in a dose of at least 300mg daily (200mg in renal impairment)  AND • Patient has not achieved a target serum uric acid level using febuxostat (Uloric) 80mg daily • Patient has a serum uric acid level above 6mg/dl despite use of Uloric • Patient is not using Zurampic as monotherapy
LIMITATIONS	Zurampic is not recommended for the treatment of asymptomatic hyperuricemia
REFERENCES	<ol> <li>Duzallo full prescribing information. Ironwood Pharmaceuticals, Inc., Cambridge, MA</li> <li>Zurampic full prescribing information. Ironwood Pharmaceuticals, Inc., Cambridge, MA</li> </ol>
P&T REVIEW HISTORY	11/16, 11/17, 1/18, 11/18
REVISION RECORD	11/17, 1/18